

Company Product	Indication & Usage	Dosage & Administration	Cost Assistance
<p><b>Glucagon</b></p> 	<p>Any age</p>	<p>Pediatric patient weighing less than 44 lb (20 kg), administer 0.5 mg intra-muscular</p> <p>Pediatric patient weighing more than 44 lb (20 kg) administer 1 mg intramuscular</p>	<p><a href="https://www.lillyglucagon.com/">https://www.lillyglucagon.com/</a></p>
<p><b>Glucagen</b></p> 	<p>Any age</p>	<p>Pediatric patient weighing less than 55 lb (25kg) administer 0.5 mL intramuscular</p> <p>Pediatric patient weighing more than 55 lb (25kg) administer 1 mL intramuscular</p>	<p><a href="https://medicineassistance.org/">https://medicineassistance.org/</a></p>
<p><b>Fresenius Kabi's Glucagon</b></p> 	<p>Any age</p>	<p>Pediatric patient weighing less than 55lb (25kg) administer 0.5mg intramuscular</p> <p>Pediatric patient weighing more than 55 lb (25kg) administer 1mg intramuscular</p>	<p><a href="https://kabicare.us/">https://kabicare.us/</a></p>
<p><b>Gvoke PFS Glucagon</b></p> 	<p>2 years and above</p>	<p>Pediatric patient who weigh less than 45kg Administer 0.5 mg subcutaneous</p> <p>Pediatric patient who weighs more than 45kg Administer 1mg subcutaneous</p>	<p><a href="https://www.gvokeglucagon.com/savings-and-support">https://www.gvokeglucagon.com/savings-and-support</a></p>
<p><b>Baqsimi</b></p> 	<p>4 years and older</p>	<p>Administer 3 mg as one actuation of the intranasal device into one nostril. If there has been no response after 15 minutes, an additional 3 mg dose of BAQSIMI from a new device may be administered while waiting for emergency assistance</p>	<p><a href="https://www.baqsimi.com/">https://www.baqsimi.com/</a></p>